ELSEVIER

Journal of Orthopaedic Research 21 (2003) 1157–1159

Journal of Orthopaedic Research

www.elsevier.com/locate/orthres

Volume 21, 2003

AUTHOR INDEX

Abbas S, 1041
Abe M, 36
Abramowitch SD, 708, 1124
Abu-Amer Y, 202, 1041
Abuzzahab F, 676
Ackermann PW, 432
Ahmed M, 465
Ahmed S, 102
Aigner T, 96
Akeson WH, 584
Akhter MP, 14
Alfredson H, 970
Algenstaedt P, 805
Almekinders L, 256
Amadio PC, 813
Amiel D, 584
Amin M, 167
Amis AA, 780
An H, 922
An K-N, 162, 813
Andersen H, 780
Anderson D, 676
Andersson GBJ, 922
Andrew G, 685
Angele P, 451
Anic T, 976
Ann Barrington N, 529
Antoniou J, 81
Anvari B, 88
Aralica G, 976
Archambault JM, 256
Arens S, 373
Aspenberg P, 196, 617
Athanasiou KA, 88
Athanasou NA, 73
Attawia MA, 1005
Aubin CÉ, 931
Augat P, 1011
Awad HA, 420
Axe MJ, 1131

Bäckman A, 970
Bäckman S, 970
Bakker FC, 521
Banes AJ, 256
Barbe MF, 167
Barr AE, 167
Barrance PJ, 1131
Batelja L, 976
Bau B, 96
Beauséjour M, 931
Beck M, 1145
Bellemans J, 123, 572
Benhaim P, 622
Bennell KL, 553, 792
Berglund LJ, 162
Bergström J, 465
Bhargava M, 888
Bishop AT, 1056
Blake CA, 997
Block JA, 1063, 1071
Blokhuis TJ, 521
Boban-Blagaic A, 976
Bogdanske J, 224
Boivin GP, 420
Bonewald LF, 326
Borovic S, 976
Boskey AL, 1
Böttcher A, 805
Bouxsein ML, 997
Boyan BD, 326, 638
Boyd A, 1049
Boyer MI, 990
Brüntrup J, 250
Bragulla H, 662
Bray RC, 787, 1118
Breme J, 213
Brodt P, 1063
Bruce WJM, 655
Buchanan TS, 1131
Buchbinder R, 792
Buckwalter J, 597
Bull AMJ, 780
Bullock-Saxton JE, 231
Bundgaard KG, 335
Bunn JR, 843
Burger J, 1011
Burghardt A, 6
Burnet S, 937
Burton-Wurster N, 730
Butler DL, 420
Bynum D, 256
Byron CD, 1025

Cain S, 655
Campbell D, 1049
Cance WG, 1071
Carandang G, 540
Chan D, 685
Chapman-Sheath PJ, 655
Cheah KSE, 685
Chen AC, 584, 872
Chen C-T, 573, 888
Chen J, 963
Chen PC, 761
Chen T-H, 1033
Chen W-Y, 365
Chern T-C, 365
Cheung KMC, 685
Chhabra A, 826
Chiba K, 922
Choi JY, 622
Choi W, 597
Christensen KS, 335
Chubinskaya S, 899
Claes L, 320, 1011
Clark RT, 826
Clemens V, 296
Clohisy JC, 202, 1041
Cochran DL, 638
Coert JH, 1151
Colwell Jr CW, 761
Conlon MA, 157
Connolly CK, 843
Costic RS, 379
Coutts RD, 584
Cowan SM, 553
Cramer H, 1090
Crawford E, 44
Crisco III JJ, 489
Crossley KM, 553, 792
Crum JA, 723
Crystal RG, 573
Cs-Szabo G, 730
Cullen DM, 14
Cullinane D, 670

Dahners LE, 183
Dai Q, 604
Dau Y-C, 365
D'Augusta D, 997
Davidson JM, 245
Davies M, 62
Dawson JM, 245
Day B, 960
Day SM, 744
Dean DD, 326, 638
De Bari C, 123, 572
Debes J, 655
de Bock C, 819
Debski RE, 379, 1107
Decker K, 914
DeHeer DH, 744
Dell'Accio F, 123, 572
Deloria LB, 881
DeLott LB, 744
den Boer FC, 521
Detwiler KB, 245
Dickson GR, 28, 843
Dimmler A, 1090
Ditsios K, 990
D'Lima DD, 761
Doi M, 36
Donahue HJ, 326
Dougall WC, 676
Dragoo JL, 622
Dresing K, 213
Dressler MR, 420
Drissi H, 908
Duck TR, 399
Duda GN, 662, 767
Dunning CE, 399, 405
Durko M, 1063
Dutton CM, 1063

Eastell R, 529, 691
Eckardt H, 335
Eckhardt C, 836
Eger W, 96
Ehrlich MG, 489
Einhorn TA, 20, 670
Eliav U, 149
Ericson M, 881
Eulert J, 1090

Farahmand F, 780
Favata M, 413
Feitelson JBA, 497
Ferris L, 352
Filler T, 250
Fink B, 805
Fischer MB, 941
Fitzsimmons JS, 296
Fleming JT, 497
Flick LM, 676
Ford JL, 54
Forslund C, 617
Foster W, 798
Frank CB, 716
Frazier E, 202
Frommlet F, 941
Frosch K-H, 213
Fukuda H, 1138
Fukuda Y, 1107
Fukunaga K, 359
Fung YK, 14

Ganz R, 1145
Gardiner JC, 1098
Gardner-Morse MG, 547
Garrett W, 256
Gaschen V, 826
Gates J, 62
Gaughan JP, 167
Gavin TM, 540
Gebhard PM, 96
Gehrsitz A, 96
Gelberman RH, 990
Georgescu H, 914
Gerstenfeld LC, 20, 670
Ghanayem AJ, 540
Gibeling JC, 481
Gibson VA, 481
Gilbert TW, 708
Gilbertson B, 752
Gillies RM, 655
Gimbel JA, 413
Giurea A, 584
Glant TT, 730
Goel V, 597
Goldberg VM, 102
Goldhahn J, 836
Goodman SB, 110, 117
Goodrich LR, 573
Goomer RS, 584
Gordon KD, 405
Gorzelany I, 167
Gotoh M, 1138
Greshake O, 250
Grodzinsky AJ, 272
Gruber R, 941
Grynpas MD, 132, 590

doi:10.1016/S0736-0266(03)00233-X

Guilak F, 963
Guyton G, 256

Haarman HJThM, 521
Haas NP, 662
Hajdarevic K, 976
Hak DJ, 503
Halder AM, 162
Hall DJ, 697
Hamada K, 1138
Hamer AJ, 529, 691
Hamrick MW, 1025
Han B, 648
Handa A, 1138
Hansen ES, 335
Hansen-Algenstaedt N, 805
Haqqi TM, 102
Hari Reddi A, 503
Hashimoto J, 44
Hashizume H, 535
Havey RM, 540
Haynatzki GR, 14
Hazelwood SJ, 481, 503
Hazleman BL, 937
Hedrick MH, 622
Heers G, 162
Heisey D, 224
Hejna MJ, 139
Heller M, 662
Heller MO, 767
Hempfing A, 1145
Hendrich C, 1090
Henke T, 1011
Henneberg M, 352
Heretis K, 899
Hermida JC, 761
Herzog J, 442
Herzog W, 256
Hesseln J, 836
Hidaka C, 573
Hiller LP, 481
Himori K, 386
Hinman RS, 792
Hirayama T, 202
Hodges PW, 553
Holbein O, 320
Hotta T, 949
Hovius SER, 1151
Howell SM, 459
Hsu C-C, 365, 984
Hsu H-P, 272
Huang A, 459
Huang C-S, 984
Huang J, 622
Huang W, 88
Huard J, 798
Huk OL, 81
Hull ML, 459
Humphreys H, 937
Hunziker EB, 290, 826
Hvid I, 335

Ignatius A, 320
Ikenoue T, 110, 117, 511
Ikoma K, 149
Imakiire A, 949
Ionescu AM, 908
Ishii Y, 1113
Issever AS, 6
Ito H, 630
Itoh M, 949
Iwamoto Y, 511
Iwaniec UT, 14
Iwasaki N, 393

Jackson DW, 470
Jadrijevic S, 976
Jari R, 379
Jepsen KJ, 451
Jiang X, 1063
Jingushi S, 511
Jo M, 62
Johnson JA, 399, 405
Johnstone B, 451
Jones J, 780
Joscheck C, 805
Jou I-M, 365
Julius C, 1090

Kälicke T, 373
Kaluarachi K, 685
Kamath RV, 730
Kandel RA, 132, 590
Kandziora F, 662
Karreth F, 941
Kaspar D, 320
Kasra M, 597
Kassi J-P, 662
Kato T, 359
Kawakami M, 535
Kawamura K, 559
Keays AC, 231
Keays SL, 231
Kikukawa K, 359
Kim H, 997
Kim YH, 1071
Kimura JH, 730
King GJW, 399, 405
Kinner B, 20
Kishida Y, 850
Kleemann RU, 767
Klein P, 662
Klein TJ, 584
Knothe Tate ML, 1018
Kofron MD, 1005
Koh JL, 565
Koike C, 805
Kokubu T, 503
Kopljar M, 976
Krebsbach PH, 859
Kreicbergs A, 432, 465
Kudo O, 73
Kuettner KE, 899
Kumano F, 922
Kumar B, 899
Kunishima DH, 470
Kunz D, 224
Kuribayashi K, 535
Kurz B, 610
Kusaka Y, 149
Kusano N, 990

Labelle H, 931
Laib A, 6
LaPrade RF, 723
Laurencin CT, 1005
LeBaron RG, 88
Leblanc CW, 489
Lee CA, 20
Lee CR, 272
Lee JW, 1071
Lee MS, 110, 117
Lee Y, 62
Leonard CA, 787, 1118
Lerner UH, 970
Lester GE, 183
Leunig M, 1145
Lewis JL, 881
Lhotka C, 189

Li G, 28, 843
Li J, 432
Li RH, 997
Li X, 908
Li XJ, 997
Li Y, 798
Li Z, 326
Lieberman JR, 62, 622
Liljenqvist U, 250
Lill CA, 836
Lin EY, 110
Lin PM, 888
Lind M, 335
Liu W, 730
Liu XC, 865
Liu Y, 326
Loh JC, 1107
Lohmann CH, 213, 326, 638
Looney RJ, 1049
Lopez MJ, 224
Lorentzon M, 970
Lou J, 44
Lu W, 685
Lundeberg T, 432
Lust G, 730
Luyten FP, 123, 572
Lyon R, 865

Majumdar S, 6
Malemud CJ, 102
Manfield L, 604
Manner PA, 697
Manske P, 44
Mansour J, 451
Markel MD, 224
Markolf KL, 177
Marsh DR, 28, 843
Martin J, 597
Martin RB, 481
Masuda K, 139, 872, 922
Mathieu PA, 931
Matsumoto F, 386
Matsumoto T, 535
McAllister DR, 177
McCauley LK, 859
McColl G, 792
McConnell J, 553
McMahon PJ, 1107
McWilliams J, 177
Meade KP, 540
Meek MF, 1151
Meier J, 865
Melhem S, 914
Mentlein R, 610
Merrihew C, 899
Metcalf BR, 792
Mielke C, 670
Miki H, 850
Mikic B, 826
Minami A, 393
Miyamoto E, 359
Miyanaga Y, 1113
Miyazawa T, 393
Mizner RL, 775
Mizuguchi J, 949
Mochida J, 1138
Moore DC, 489
Morales TI, 290
Muhr G, 373
Mukherjee N, 296
Müller I, 805
Müller R, 489
Mura N, 162
Muramatsu K, 1056

Murray MM, 238, 245
Murrell GAC, 604, 819
Mushipe M, 843

Nagai S, 949
Nakajima M, 559
Nakamura M, 1138
Nakanishi K, 850
Nanney LB, 245
Nasser P, 1018
Navon G, 149
Neidlinger-Wilke C, 320
Nerlich M, 451
Newcombe P, 231
Neys J, 123, 572
Ni Q, 312
Nichols D, 937
Nimni ME, 648
Nishi H, 535
Nishii T, 850
Nixon AJ, 573
Norcross JP, 183
Nöth U, 1090
Nötzli HP, 1145
Nuber GW, 565
Nylander I, 465

Oakes DA, 177
Ochi T, 850
Odermatt D, 931
O'Driscoll SW, 162, 296
Oegema Jr TR, 881
Ohtsuka H, 36
Oizumi N, 393
Okazaki K, 511
O'Keefe RJ, 676, 908, 1049
Opsitnick NC, 1005
Orimo H, 630
Oyen-Tiesma M, 881

Pacicca DM, 20
Pang J-HS, 265
Papageorgiou CD, 708
Park HB, 1071
Patel V, 6
Patka P, 521
Patrlj L, 976
Patwardhan AG, 540
Paxinos O, 540
Pennington C, 1025
Peraus M, 1011
Perovic D, 976
Petersen W, 610
Petit A, 81
Phippard D, 670
Pilliar RM, 132, 590
Pötzl W, 250
Pope R, 352
Prater CS, 481
Press JM, 565
Printzen G, 373
Pufe T, 610
Puzas JE, 638, 908

Qi W-n, 282, 1063

Rabba J, 213
Raiss RX, 1081
Rayan V, 305
Recker RR, 14
Reddy GK, 738
Reinholz GG, 296
Rice K, 238
Richards C, 937

Ries M, 6
Ritty TM, 442
Roberts CS, 497
Robinson DE, 54
Rodosky MW, 379
Rogers A, 691
Rompe JD, 958
Rosier RN, 908
Rowell PP, 497
Rudy T, 213
Rueger D, 617
Rueger DC, 899
Rühli FJ, 352
Russell LC, 752
Rüther W, 805

Sabokbar A, 73
Safadi FF, 167
Sah RL, 139, 584, 872
Sakai H, 511
Sakai T, 341, 850
Salo PT, 787, 1118
Sanyal A, 296
Sarkar G, 296
Scammell BE, 54
Schütze N, 1090
Schaefer C, 805
Schaffler MB, 1018
Schell H, 662
Schild D, 213
Schlegel U, 373, 836
Schmid T, 922
Schneider A, 859
Schneider E, 373, 836
Schorlemmer S, 1011
Schurman DJ, 110
Schurman DJ, 117
Schwartz Z, 326, 638
Schwarz E, 62
Schwarz EM, 676, 908, 1049
Schwarzloh B, 805
Scully SP, 282, 1063, 1071
Sebecic B, 976
Seeherman HJ, 997
Seibert K, 670
Seiwerth S, 976
Senavongse W, 780
Seo Y, 149
Setton LA, 963
Sharkey PF, 697
Sheehan KM, 744
Shih H-N, 1033
Shih L-Y, 1033
Shimada T, 630
Shrive NG, 716
Sikiric P, 976
Silva MJ, 990
Simon BJ, 326
Simon TM, 470
Simonds J, 540
Simpson H, 28
Singh R, 102
Skoglund B, 196
Smith C, 451
Smith FNL, 420
Smith M, 792
Smith RL, 110, 117
Snijders CJ, 1151
Snyder-Mackler L, 775, 1131
Söder S, 96
Solomon LB, 352
Somogyi G, 798
Sondergeld I, 213
Soslowsky LJ, 413
Spector M, 20, 238, 272
Speed CA, 937, 961
Spindler KP, 245
Spiteri CG, 590
Srdjak M, 976
Staresinic M, 976
Steffan I, 189
Steinbeck J, 250
Steinert A, 1090
Steinmeyer J, 1081
Stevens JE, 775
Stockley I, 529, 691
Stoeckle U, 767
Stokes IA, 547
Stover SM, 481
Streitparth F, 662
Studer RK, 914
Stuermer KM, 213
Suenaga N, 393
Sugano N, 341, 850
Sugiyama O, 630
Suknaic S, 976
Sumner DR, 305, 730
Suzuki H, 949
Suzuki S, 630
Svagr B, 670
Svehla MJ, 655
Sylvia VL, 326, 638
Szekeres T, 189

Taboas JM, 859
Takada E, 949
Takagi K, 359
Takamiya H, 149
Takeda I, 559
Takegami K, 922
Tamaki T, 535
Tami AE, 1018
Tanaka T, 813
Tang B, 648
Tang F-T, 265
Tang P, 1107
Tarter JT, 245
Tashman S, 730
Tateishi T, 1113
Taylor J, 352
Taylor WR, 767
Thiede M, 670
Thomopoulos S, 413
Thompson Jr RC, 881
Thonar E, 922
Thonar EJ-MA, 139, 305, 872
Thornton GM, 716
Tillmann B, 610
Tomas FM, 157
Torres JH, 88
Torzilli PA, 888
Totterman S, 1049
Trindade MCD, 110, 117
Trudel G, 386
Tsai W-C, 265
Tsokos M, 610
Tsou D, 826
Tsuchida H, 44
Tsuda E, 1107, 1124
Tsuji T, 341
Tsuzaki M, 256
Tuan RS, 697
Tuli R, 697
Turcic I, 976

Ueyama Y, 1138
Uhthoff HK, 386
Ulrich-Vinther M, 676
Upton ML, 963
Urabe K, 511
Usas A, 798
Ushida T, 1113

Valenzuela RG, 1056
van der Meulen MCH, 1
van Dijke GAH, 1151
Van Sickle DC, 470
Vanderby Jr R, 224
Vanlauwe J, 123, 572
Voronov LI, 540

Waldman SD, 132, 590
Walsh WR, 655
Wang C-J, 984
Wang F-S, 984
Wang G, 565
Wang ML, 697
Wang S-T, 597
Wang X, 312
Wang Y, 604, 819
Warren RF, 573
Watzek G, 941
Weaver JM, 676
Weber M, 1090
Weinhold P, 183
Weiss JA, 1098
Weng L-H, 984
Wenner K, 752
Wentorf FA, 723
Wies J, 937
Wilkinson JM, 529, 691
Williams GN, 1131
Williams GR, 413
Williams JM, 305
Williamson AK, 872
Wippermann BW, 521
Withrow JD, 708
Wolf A, 1081
Wong M-K, 265
Wong N, 117
Woo SL-Y, 708, 1107, 1124
Wozney JM, 997
Wright RJ, 238
Wright TM, 1
Wu J, 62

Xia W, 819
Xu L-H, 1071

Yagi M, 1124
Yamaga M, 359
Yamamoto K, 949
Yamashita K, 630
Yamazaki H, 1138
Yanagisawa K, 1138
Yang KD, 984
Yang L-C, 984
Yang X, 256
Yeh OC, 481
Yokota A, 36
Yoo JU, 451
Yoshikawa H, 341, 850
Young CR, 177
Young RG, 420

Zarkovic N, 976
Zhang J, 622
Zhang L-Q, 565
Zhang X, 62, 676, 908
Zhao C, 162, 813
Zhuber K, 189
Zimmermann U, 1090
Zobitz ME, 162, 813
Zuk PA, 622
Zukor DJ, 81
Zuscik MJ, 908
Zweymüller K, 189

ELSEVIER

Journal of Orthopaedic Research 21 (2003) 1160–1171

Journal of Orthopaedic Research

www.elsevier.com/locate/orthres

Volume 21, 2003

SUBJECT INDEX

Achilles tendon
Evaluation of collagen fiber maturation and ordering in regenerating tendons employing H-1 double quantum filtered NMR spectroscopy, 149

Achilles tendon transection
Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth, 976

ACL injury
Electromyographic analysis of a modified maneuver for quadriceps femoris muscle setting with co-contraction of the hamstrings, 559

Adenovirus
Bone induction by *BMP-2* transduced stem cells derived from human fat, 622

Adhesion
The C-terminal domain of focal adhesion kinase reduces the tumor cell invasiveness in chondrosarcoma cell lines, 1071

Adhesiveness
Temporal effects of cell adhesion on mechanical characteristics of the single chondrocyte, 88

Adipose-derived stem cell
Bone induction by *BMP-2* transduced stem cells derived from human fat, 622

Adolescent idiopathic scoliosis
Electromyography of scoliotic patients treated with a brace, 931

Aggrecan
Spontaneous and experimental osteoarthritis in dog: similarities and differences in proteoglycan levels, 730

Aging
Quadriceps strength and volitional activation before and after total knee arthroplasty for osteoarthritis, 775

Alginate
Chondrogenic differentiation of mesenchymal progenitor cells encapsulated in ultrahigh-viscosity alginate, 1090

Alkaline phosphatase
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Allotransplantation
Detection of chimerism following vascularized bone allotransplantation by polymerase chain reaction using a Y-chromosome specific primer, 1056

ALP, alkaline phosphatase
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Alpha-smooth muscle actin
Inhibition of tendon cell migration by dexamethasone is correlated with reduced alpha-smooth muscle actin gene expression: a potential mechanism of delayed tendon healing, 265

Angiogenesis
Bilaterally increased VEGF-levels in muscles during experimental unilateral callus distraction, 805
Correlation of healing capacity with vascular response in the anterior cruciate and medial collateral ligaments of the rabbit, 1118
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335
Mechanical factors influence the expression of endostatin—an inhibitor of angiogenesis—in tendons, 610
Physiologic weight-bearing increases new vessel formation during distraction osteogenesis: a micro-tomographic imaging study, 489
Shock wave therapy induces neovascularization at the tendon–bone junction A study in rabbits, 984
Vascular adaptation of intact joint stabilizing structures in the posterior cruciate ligament deficient rabbit knee, 787

Animal model
Biomechanical evaluation of healing in a non-critical defect in a large animal model of osteoporosis, 836
Development of an atrophic nonunion model and comparison to a closed healing fracture in rat femur, 503
Influence of a standardized closed soft tissue trauma on resistance to local infection. An experimental study in rats, 373
Shear movement at the fracture site delays healing in a diaphyseal fracture model, 1011

Animal movement
A reliable externally fixated murine femoral fracture model that accounts for variation in movement between animals, 843

Announcement
760
First Announcement: Japanese–Korean Joint Conference on Rehabilitation Medicine 2004, 962

Anterior cruciate ligament
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224
Correlation of healing capacity with vascular response in the anterior cruciate and medial collateral ligaments of the rabbit, 1118
Specificity of muscle action after anterior cruciate ligament injury, 1131
Spontaneous and experimental osteoarthritis in dog: similarities and differences in proteoglycan levels, 730
The healing medial collateral ligament following a combined anterior cruciate and medial collateral ligament injury—a biomechanical study in a goat model, 1124
The relationship between knee strength and functional stability before and after anterior cruciate ligament reconstruction, 231
Vascular adaptation of intact joint stabilizing structures in the posterior cruciate ligament deficient rabbit knee, 787

Anteversion
THA loading arising from increased femoral anteversion and offset may lead to critical cement stresses, 767

Anti-angiogenic
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Antisense RNA
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

Apoptosis
Chemotherapeutic agents sensitize sarcoma cell lines to tumor necrosis factor-related apoptosis-inducing ligand-induced caspase-8 activation, apoptosis and loss of mitochondrial membrane potential, 949
Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles, 697
Effect of pamidronate on the stimulation of macrophage TNF-α release by ultra-high-molecular-weight polyethylene particles: a role for apoptosis, 81

Hyalgan® has a dose-dependent differential effect on macrophage proliferation and cell death, 744
Rapid new bone tissue remodeling during distraction osteogenesis is associated with apoptosis, 28

Area
Measurement of lesion area and volume by three-dimensional spoiled gradient-echo MR imaging in osteonecrosis of the femoral head, 850

Arteriolar blockade
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Arthritis
Epigallocatechin-3-gallate selectively inhibits interleukin-1β-induced activation of mitogen activated protein kinase subgroup c-*Jun N*-terminal kinase in human osteoarthritis chondrocytes, 102
Nitric oxide inhibition of IGF-1 stimulated proteoglycan synthesis: role of cGMP, 914

Articular cartilage
Long-term intermittent shear deformation improves the quality of cartilaginous tissue formed in vitro, 590
Tensile mechanical properties of bovine articular cartilage: variations with growth and relationships to collagen network components, 872
The use of specific chondrocyte populations to modulate the properties of tissue-engineered cartilage, 132
Time, stress, and location dependent chondrocyte death and collagen damage in cyclically loaded articular cartilage, 888

Articular cartilage repair
The use of intra-articular Na-hyaluronate as a potential chondroprotective device in experimentally induced acute articular cartilage injury and repair in rabbits, 305

Articular chondrocytes
Type II collagen modulates the composition of extracellular matrix synthesized by articular chondrocytes, 282

Aseptic loosening
Bone mineral density and biochemical markers of bone turnover in aseptic loosening after total hip arthroplasty, 691

Autograft
The level of compressive load affects conclusions from statistical analyses to determine whether a lateral meniscal autograft restores tibial contact pressure to normal: a study in human cadaveric knees, 459

Autologous osteoblasts
Autologous osteoblasts enhance osseointegration of porous titanium implants, 213

Axial compressive preload
Physiological axial compressive preloads increase motion segment stiffness, linearity and hysteresis in all six degrees of freedom for small displacements about the neutral posture, 547

Basic fibroblast growth factor
Platelets are mitogenic for periosteum-derived cells, 941

Biceps tendon
Gliding properties of the long head of the biceps brachii, 162

Bicipital groove
Gliding properties of the long head of the biceps brachii, 162

Bioassay
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Biochemical markers of bone turnover
Bone mineral density and biochemical markers of bone turnover in aseptic loosening after total hip arthroplasty, 691

Biodegradable polymer
Cryopreservation of tissue engineered constructs for bone, 1005

Biomechanical
Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth, 976

Biomechanical properties
The healing medial collateral ligament following a combined anterior cruciate and medial collateral ligament injury—a biomechanical study in a goat model, 1124

Biomechanics
Adhesion of perichondrial cells to a polylactic acid scaffold, 584
Biomechanical evaluation of healing in a non-critical defect in a large animal model of osteoporosis, 836
Fibronectin metabolism of cartilage explants in response to the frequency of intermittent loading, 1081
Physiological axial compressive preloads increase motion segment stiffness, linearity and hysteresis in all six degrees of freedom for small displacements about the neutral posture, 547
Shear movement at the fracture site delays healing in a diaphyseal fracture model, 1011
Tensile mechanical properties of bovine articular cartilage: variations with growth and relationships to collagen network components, 872

Biphosphonate
Effect of pamidronate on the stimulation of macrophage TNF-α release by ultra-high-molecular-weight polyethylene particles: a role for apoptosis, 81

Blood flow
Correlation of healing capacity with vascular response in the anterior cruciate and medial collateral ligaments of the rabbit, 1118
Vascular adaptation of intact joint stabilizing structures in the posterior cruciate ligament deficient rabbit knee, 787

Blood levels
Four-year study of cobalt and chromium blood levels in patients managed with two different metal-on-metal total hip replacements, 189

BMD
Bone loss following tendon laceration, repair and passive mobilization, 990

BMP
Chondrogenic differentiation of mesenchymal progenitor cells encapsulated in ultrahigh-viscosity alginate, 1090

BMP-2
Bone induction by *BMP-2* transduced stem cells derived from human fat, 622
rhBMP-2 injected in a calcium phosphate paste (α-BSM) accelerates healing in the rabbit ulnar osteotomy model, 997

BMP-2, bone morphogenic protein-2
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

BMP-3, bone morphogenic protein-3 (osteogenin)
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

BMPs, bone morphogenetic proteins
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Bone
Bone induction by *BMP-2* transduced stem cells derived from human fat, 622
Determination of cortical bone porosity and pore size distribution using a low field pulsed NMR approach, 312
MicroCT evaluation of normal and osteoarthritic bone structure in human knee specimens, 6
Noninvasive fatigue fracture model of the rat ulna, 1018
Purification and quantification of opioid peptides in bone and joint tissues—a methodological study in the rat, 465

Bone blood flow
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Bone formation
Skeletal homeostasis in tissue-engineered bone, 859

Bone ingrowth
The effects of sex and estrogen therapy on bone ingrowth into porous coated implant, 1033

Bone lengthening
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335
Physiologic weight-bearing increases new vessel formation during distraction osteogenesis: a micro-tomographic imaging study, 489

Bone marrow
Cyclic hydrostatic pressure enhances the chondrogenic phenotype of human mesenchymal progenitor cells differentiated in vitro, 451

Bone marrow stromal cells
Skeletal homeostasis in tissue-engineered bone, 859

Bone mineral content (BMC)
Effects of nicotine on bone mass and strength in aged female rats, 14

Bone mineral density
Bone mineral density and biochemical markers of bone turnover in aseptic loosening after total hip arthroplasty, 691
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Bone mineral density (BMD)
Effects of nicotine on bone mass and strength in aged female rats, 14

Bone morphogenetic protein
Smad6 is induced by BMP-2 and modulates chondrocyte differentiation, 908

Bone morphogenetic protein-7
Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads, 922

Bone morphogenetic proteins
Alterations in endogenous osteogenic protein-1 with degeneration of human articular cartilage, 899
GDF-5 deficiency in mice delays Achilles tendon healing, 826

Bone regeneration
Platelets are mitogenic for periosteum-derived cells, 941

Bone remodeling
Rapid new bone tissue remodeling during distraction osteogenesis is associated with apoptosis, 28

Bone repair
The fate of soft callus chondrocytes during long bone fracture repair, 54

Bone resorption
Bone loss following tendon laceration, repair and passive mobilization, 990
Skeletal homeostasis in tissue-engineered bone, 859

Bone strength
Bone architecture and disc degeneration in the lumbar spine of mice lacking GDF-8 (myostatin), 1025
Effects of nicotine on bone mass and strength in aged female rats, 14

Bone wafers
Pretreatment of bone with osteoclasts affects phenotypic expression of osteoblast-like cells, 638

Boston brace
Electromyography of scoliotic patients treated with a brace, 931

Bound water
Evaluation of collagen fiber maturation and ordering in regenerating tendons employing H-1 double quantum filtered NMR spectroscopy, 149

C2C12 cells
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Calcium polyphosphate substrates
Long-term intermittent shear deformation improves the quality of cartilaginous tissue formed in vitro, 590
The use of specific chondrocyte populations to modulate the properties of tissue-engineered cartilage, 132

Callostasis
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Callus distraction
Bilaterally increased VEGF-levels in muscles during experimental unilateral callus distraction, 805

Canine
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224

Carrier
rhBMP-2 injected in a calcium phosphate paste (α-BSM) accelerates healing in the rabbit ulnar osteotomy model, 997

Cartilage
Cell death after cartilage impact occurs around matrix cracks, 881
Expanded phenotypically stable chondrocytes persist in the repair tissue and contribute to cartilage matrix formation and structural integration in a goat model of autologous chondrocyte implantation, 123
Fibronectin metabolism of cartilage explants in response to the frequency of intermittent loading, 1081
Nitric oxide inhibition of IGF-1 stimulated proteoglycan synthesis: role of cGMP, 914
Spontaneous and experimental osteoarthritis in dog: similarities and differences in proteoglycan levels, 730
The fate of soft callus chondrocytes during long bone fracture repair, 54

Cartilage biology
Induction of CD-RAP mRNA during periosteal chondrogenesis, 296

Cartilage degeneration
Alterations in endogenous osteogenic protein-1 with degeneration of human articular cartilage, 899

Cartilage repair
Adhesion of perichondrial cells to a polylactic acid scaffold, 584

Cartilage-derived retinoic acid-sensitive protein
Induction of CD-RAP mRNA during periosteal chondrogenesis, 296

Cell adhesion
Adhesion of perichondrial cells to a polylactic acid scaffold, 584
Temporal effects of cell adhesion on mechanical characteristics of the single chondrocyte, 88

Cell death
The fate of soft callus chondrocytes during long bone fracture repair, 54
Time, stress, and location dependent chondrocyte death and collagen damage in cyclically loaded articular cartilage, 888

Cell migration
Inhibition of cell migration from tendon explants into fibrin clots by extracts derived from cheese whey is largely due to transforming growth factor-beta, 157
Inhibition of tendon cell migration by dexamethasone is correlated with reduced alpha-smooth muscle actin gene expression: a potential mechanism of delayed tendon healing, 265

Cell viability
Cell death after cartilage impact occurs around matrix cracks, 881

Cells
Effect of dynamic hydrostatic pressure on rabbit intervertebral disc cells, 597

cGMP dependent kinase
Nitric oxide inhibition of IGF-1 stimulated proteoglycan synthesis: role of cGMP, 914

Chemotherapy
Chemotherapeutic agents sensitize sarcoma cell lines to tumor necrosis factor-related apoptosis-inducing ligand-induced caspase-8 activation, apoptosis and loss of mitochondrial membrane potential, 949

Chimerism
Detection of chimerism following vascularized bone allotransplantation by polymerase chain reaction using a Y-chromosome specific primer, 1056

Chondrocyte
Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes, 96

Smad6 is induced by BMP-2 and modulates chondrocyte differentiation, 908

Chondrocyte labeling
Expanded phenotypically stable chondrocytes persist in the repair tissue and contribute to cartilage matrix formation and structural integration in a goat model of autologous chondrocyte implantation, 123

Chondrocytes
Long-term intermittent shear deformation improves the quality of cartilaginous tissue formed in vitro, 590
Nitric oxide inhibition of IGF-1 stimulated proteoglycan synthesis: role of cGMP, 914
The use of specific chondrocyte populations to modulate the properties of tissue-engineered cartilage, 132

Chondrocytes/*transplantation
Expanded phenotypically stable chondrocytes persist in the repair tissue and contribute to cartilage matrix formation and structural integration in a goat model of autologous chondrocyte implantation, 123

Chondrogenesis
Cyclic hydrostatic pressure enhances the chondrogenic phenotype of human mesenchymal progenitor cells differentiated in vitro, 451
Induction of CD-RAP mRNA during periosteal chondrogenesis, 296

Chondrogenic differentiation
Chondrogenic differentiation of mesenchymal progenitor cells encapsulated in ultrahigh-viscosity alginate, 1090

Chondrosarcoma
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063
The C-terminal domain of focal adhesion kinase reduces the tumor cell invasiveness in chondrosarcoma cell lines, 1071

Chromium
Four-year study of cobalt and chromium blood levels in patients managed with two different metal-on-metal total hip replacements, 189

Closed soft tissue trauma
Influence of a standardized closed soft tissue trauma on resistance to local infection. An experimental study in rats, 373

Co-contraction
Electromyographic analysis of a modified maneuver for quadriceps femoris muscle setting with co-contraction of the hamstrings, 559

Cobalt
Four-year study of cobalt and chromium blood levels in patients managed with two different metal-on-metal total hip replacements, 189

Collagen
Differential effects of static and dynamic compression on meniscal cell gene expression, 963
Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes, 96
Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads, 922
Tensile mechanical properties of bovine articular cartilage: variations with growth and relationships to collagen network components, 872
Type II collagen modulates the composition of extracellular matrix synthesized by articular chondrocytes, 282

Collagen damage
Time, stress, and location dependent chondrocyte death and collagen damage in cyclically loaded articular cartilage, 888

Compression
Differential effects of static and dynamic compression on meniscal cell gene expression, 963
The level of compressive load affects conclusions from statistical analyses to determine whether a lateral meniscal autograft restores tibial contact pressure to normal: a study in human cadaveric knees, 459

Connective tissue matrix
Glucose-mediated in vitro glycation modulates biomechanical integrity of the soft tissues but not hard tissues, 738

Connexin 43
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Contact pressure
The level of compressive load affects conclusions from statistical analyses to determine whether a lateral meniscal autograft restores tibial contact pressure to normal: a study in human cadaveric knees, 459

Cortical porosity
Determination of cortical bone porosity and pore size distribution using a low field pulsed NMR approach, 312

Cox enzymes
Differential inhibition of fracture healing by non-selective and cyclooxygenase-2 selective non-steroidal anti-inflammatory drugs, 670

Cox-2
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Crack
Cell death after cartilage impact occurs around matrix cracks, 881

Cross-linking
Tensile mechanical properties of bovine articular cartilage: variations with growth and relationships to collagen network components, 872

Cryopreservation
Cryopreservation of tissue engineered constructs for bone, 1005

Cytodetacher
Temporal effects of cell adhesion on mechanical characteristics of the single chondrocyte, 88

Cytokine
Epigallocatechin-3-gallate selectively inhibits interleukin-1β-induced activation of mitogen activated protein kinase subgroup c-*Jun N*-terminal kinase in human osteoarthritis chondrocytes, 102

Cytokines
Differences in the cytokine profiles associated with prostate cancer cell induced osteoblastic and osteolytic lesions in bone, 62
IL-1β induces COX2, MMP-1, -3 and -13, ADAMTS-4, IL-1β and IL-6 in human tendon cells, 256

DBM, demineralized bone matrix
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Decorin
Differential effects of static and dynamic compression on meniscal cell gene expression, 963

Demineralized bone matrix
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Dexamethasone
Inhibition of tendon cell migration by dexamethasone is correlated with reduced alpha-smooth muscle actin gene expression: a potential mechanism of delayed tendon healing, 265

Differentiation
Pretreatment of bone with osteoclasts affects phenotypic expression of osteoblast-like cells, 638
Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes, 96

Disability
Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis, 792

Dislocation
Quantitative measurement of patellofemoral joint stability: force–displacement behavior of the human patella in vitro, 780

Distraction osteogenesis
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Physiologic weight-bearing increases new vessel formation during distraction osteogenesis: a micro-tomographic imaging study, 489
Rapid new bone tissue remodeling during distraction osteogenesis is associated with apoptosis, 28

DQF MRS
Evaluation of collagen fiber maturation and ordering in regenerating tendons employing H-1 double quantum filtered NMR spectroscopy, 149

dsRNA, double-stranded RNA
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

Dynamic
Effect of dynamic hydrostatic pressure on rabbit intervertebral disc cells, 597

Editorial
Guidelines for describing mouse skeletal phenotype, 1

EGF, epithelial growth factor
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Electromyography
Electromyography of scoliotic patients treated with a brace, 931
Simultaneous feedforward recruitment of the vasti in untrained postural tasks can be restored by physical therapy, 553

EMG
Electromyographic analysis of a modified maneuver for quadriceps femoris muscle setting with co-contraction of the hamstrings, 559
Specificity of muscle action after anterior cruciate ligament injury, 1131

End-load bending
Noninvasive fatigue fracture model of the rat ulna, 1018

Endochondral ossification
Smad6 is induced by BMP-2 and modulates chondrocyte differentiation, 908

eNOS
Shock wave therapy induces neovascularization at the tendon–bone junction A study in rabbits, 984

ERKs
Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts, 1041

Erratum
572, 759

Estrogen replacement therapy
The effects of sex and estrogen therapy on bone ingrowth into porous coated implant, 1033

Experimental arthritis
The use of intra-articular Na-hyaluronate as a potential chondroprotective device in experimentally induced acute articular cartilage injury and repair in rabbits, 305

External fixation
A reliable externally fixated murine femoral fracture model that accounts for variation in movement between animals, 843
An externally fixed femoral fracture model for mice, 685

External fixator
The initial phase of fracture healing is specifically sensitive to mechanical conditions, 662

Extracellular matrix
Adhesion of perichondrial cells to a polylactic acid scaffold, 584
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819
Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads, 922
Tendon cells produce gelatinases in response to type I collagen attachment, 442

Fat accumulation
Early reattachment does not reverse atrophy and fat accumulation of the supraspinatus—an experimental study in rabbits, 386

Fatigue
Osteon pullout in the equine third metacarpal bone: effects of ex vivo fatigue, 481

Fatigue loading
Noninvasive fatigue fracture model of the rat ulna, 1018

Femoral head
Measurement of lesion area and volume by three-dimensional spoiled gradient-echo MR imaging in osteonecrosis of the femoral head, 850

Femoral perfusion
Evaluation of femoral perfusion in a nontraumatic rabbit osteonecrosis model with T2'-weighted dynamic MRI, 341

Femur
An externally fixed femoral fracture model for mice, 685

FGF, fibroblast growth factor
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Fibrocartilage
Mechanical factors influence the expression of endostatin—an inhibitor of angiogenesis—in tendons, 610

Fibrocartilage reformation
Early reattachment does not reverse atrophy and fat accumulation of the supraspinatus—an experimental study in rabbits, 386

Fibronectin
Fibronectin metabolism of cartilage explants in response to the frequency of intermittent loading, 1081

Fibrosis
Gamma interferon as an antifibrosis agent in skeletal muscle, 798

Fibular collateral ligament
The anatomy of the posterolateral aspect of the rabbit knee, 723

Finite element
Subject-specific finite element analysis of the human medial collateral ligament during valgus knee loading, 1098

Finite element analysis
THA loading arising from increased femoral anteversion and offset may lead to critical cement stresses, 767

Fixation
The level of compressive load affects conclusions from statistical analyses to determine whether a lateral meniscal autograft restores tibial contact pressure to normal: a study in human cadaveric knees, 459

Flexion–extension
Effect of compressive follower preload on the flexion–extension response of the human lumbar spine, 540

Focal adhesion kinase
The C-terminal domain of focal adhesion kinase reduces the tumor cell invasiveness in chondrosarcoma cell lines, 1071

Follower load
Effect of compressive follower preload on the flexion–extension response of the human lumbar spine, 540

Foot
Extracorporeal shock wave therapy for plantar fasciitis. A double blind randomised controlled trial, 937

Fracture
An externally fixed femoral fracture model for mice, 685
Expression of parathyroid hormone-related peptide and insulin-like growth factor I during rat fracture healing, 511
Osteon pullout in the equine third metacarpal bone: effects of ex vivo fatigue, 481
rhBMP-2 injected in a calcium phosphate paste (α-BSM) accelerates healing in the rabbit ulnar osteotomy model, 997

Fracture healing
A reliable externally fixated murine femoral fracture model that accounts for variation in movement between animals, 843
Biomechanical evaluation of healing in a non-critical defect in a large animal model of osteoporosis, 836
Development of an atrophic nonunion model and comparison to a closed healing fracture in rat femur, 503
Effects of receptor activator of NFκB (RANK) signaling blockade on fracture healing, 676

Noninvasive fatigue fracture model of the rat ulna, 1018
The initial phase of fracture healing is specifically sensitive to mechanical conditions, 662

Fracture repair
Differential inhibition of fracture healing by non-selective and cyclooxygenase-2 selective non-steroidal anti-inflammatory drugs, 670
Shear movement at the fracture site delays healing in a diaphyseal fracture model, 1011

Friction
Gliding properties of the long head of the biceps brachii, 162

Function
The relationship between knee strength and functional stability before and after anterior cruciate ligament reconstruction, 231

G418, geneticin
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

Gadopentetate dimeglumine
Evaluation of femoral perfusion in a non-traumatic rabbit osteonecrosis model with $T2^*$-weighted dynamic MRI, 341

Gamma interferon
Gamma interferon as an antifibrosis agent in skeletal muscle, 798

Gap junctions
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Gastric pentadecapeptide BPC 157 functional
Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth, 976

GDF-5
GDF-5 deficiency in mice delays Achilles tendon healing, 826

Gelatinase
Tendon cells produce gelatinases in response to type I collagen attachment, 442

Gender
The effects of sex and estrogen therapy on bone ingrowth into porous coated implant, 1033

Gene expression
Differential effects of static and dynamic compression on meniscal cell gene expression, 963
Induction of CD-RAP mRNA during periosteal chondrogenesis, 296
Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes, 96

Gliding tendon
Mechanical factors influence the expression of endostatin—an inhibitor of angiogenesis—in tendons, 610

Goat
The healing medial collateral ligament following a combined anterior cruciate and medial collateral ligament injury—a biomechanical study in a goat model, 1124

Green tea
Epigallocatechin-3-gallate selectively inhibits interleukin-1β-induced activation of mitogen activated protein kinase subgroup c-*Jun N*-terminal kinase in human osteoarthritis chondrocytes, 102

Growth
Tensile mechanical properties of bovine articular cartilage: variations with growth and relationships to collagen network components, 872

Growth of cultured rat tendocytes
Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth, 976

Growth plate
The effects of therapeutic vs. high-intensity ultrasound on the rabbit growth plate, 865

Growth/differentiation factors
GDF-5 deficiency in mice delays Achilles tendon healing, 826

Hamstrings
Electromyographic analysis of a modified maneuver for quadriceps femoris muscle setting with co-contraction of the hamstrings, 559

Healing
The healing medial collateral ligament following a combined anterior cruciate and medial collateral ligament injury—a biomechanical study in a goat model, 1124

Heel
Extracorporeal shock wave therapy for plantar fasciitis. A double blind randomised controlled trial, 937

Hip arthroplasty
Effect of pamidronate on the stimulation of macrophage TNF-α release by ultra-high-molecular-weight polyethylene particles: a role for apoptosis, 81

Histology
Shear movement at the fracture site delays healing in a diaphyseal fracture model, 1011
The effects of therapeutic vs. high-intensity ultrasound on the rabbit growth plate, 865

Histomorphometry
A reliable externally fixated murine femoral fracture model that accounts for variation in movement between animals, 843
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Horse
Osteon pullout in the equine third metacarpal bone: effects of ex vivo fatigue, 481

Human articular cartilage
Alterations in endogenous osteogenic protein-1 with degeneration of human articular cartilage, 899

Hyalgan®
Hyalgan® has a dose-dependent differential effect on macrophage proliferation and cell death, 744

Hyaluronic acid
Hyalgan® has a dose-dependent differential effect on macrophage proliferation and cell death, 744
The use of intra-articular Na-hyaluronate as a potential chondroprotective device in experimentally induced acute articular cartilage injury and repair in rabbits, 305

Hydrostatic pressure
Effect of dynamic hydrostatic pressure on rabbit intervertebral disc cells, 597

Hysteresis
Physiological axial compressive preloads increase motion segment stiffness, linearity and hysteresis in all six degrees of freedom for small displacements about the neutral posture, 547

iaDBM, guanidine inactivated DBM
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

IGF, insulin-like growth factor
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

IL-1β
IL-1β induces COX2, MMP-1, -3 and -13, ADAMTS-4, IL-1β and IL-6 in human tendon cells, 256

Immobilization
Bone loss following tendon laceration, repair and passive mobilization, 990

Immunohistochemistry
Expression of parathyroid hormone-related peptide and insulin-like growth factor I during rat fracture healing, 511

Impact
Cell death after cartilage impact occurs around matrix cracks, 881

Implant particles
RANKL is an essential cytokine mediator of polymethylmethacrylate particle-induced osteoclastogenesis, 202

In situ hybridization
Expression of parathyroid hormone-related peptide and insulin-like growth factor I during rat fracture healing, 511

In vivo
Expanded phenotypically stable chondrocytes persist in the repair tissue and contribute to cartilage matrix formation and structural integration in a goat model of autologous chondrocyte implantation, 123
In vivo load sharing among the quadriceps components, 565

In vivo stiffness
Biomechanical evaluation of healing in a non-critical defect in a large animal model of osteoporosis, 836

Infection
Influence of a standardized closed soft tissue trauma on resistance to local infection. An experimental study in rats, 373

Inflammation
Epigallocatechin-3-gallate selectively inhibits interleukin-1β-induced activation of mitogen activated protein kinase subgroup c-*Jun N*-terminal kinase in human osteoarthritis chondrocytes, 102

Insertion site
Variation of biomechanical, structural, and compositional properties along the tendon to bone insertion site, 413

Instability
Quantitative measurement of patellofemoral joint stability: force–displacement behavior of the human patella in vitro, 780

Insulin-like growth factor I
Expression of parathyroid hormone-related peptide and insulin-like growth factor I during rat fracture healing, 511

Interfragmentary movements
The initial phase of fracture healing is specifically sensitive to mechanical conditions, 662

Intervertebral disc
Effect of dynamic hydrostatic pressure on rabbit intervertebral disc cells, 597
Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads, 922

Invasion
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063
The C-terminal domain of focal adhesion kinase reduces the tumor cell invasiveness in chondrosarcoma cell lines, 1071

Isometric tasks
Electromyography of scoliotic patients treated with a brace, 931

JJ012, a human chondrosarcoma cell line
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

JNK
Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts, 1041

Joint geometry
THA loading arising from increased femoral anteversion and offset may lead to critical cement stresses, 767

Joint laxity
Vascular adaptation of intact joint stabilizing structures in the posterior cruciate ligament deficient rabbit knee, 787

Joint mechanics
Subject-specific finite element analysis of the human medial collateral ligament during valgus knee loading, 1098

Joint stability
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224

Joints
Purification and quantification of opioid peptides in bone and joint tissues—a methodological study in the rat, 465

Kinematics
Effect of compressive follower preload on the flexion–extension response of the human lumbar spine, 540

Knee
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224
A quantitative analysis of valgus torque on the ACL: a human cadaveric study, 1107
In vivo load sharing among the quadriceps components, 565
MicroCT evaluation of normal and osteoarthritic bone structure in human knee specimens, 6
Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis, 792

Knee anatomy
The anatomy of the posterolateral aspect of the rabbit knee, 723

Knee arthroplasty
Quadriceps strength and volitional activation before and after total knee arthroplasty for osteoarthritis, 775

Knee stability
Specificity of muscle action after anterior cruciate ligament injury, 1131

Laxity
Quantitative measurement of patellofemoral joint stability: force–displacement behavior of the human patella in vitro, 780

Letters to the Editor
958, 960

Ligament
Effects of transcutaneous topical injection of oxygen on vascular endothelial growth factor gene into the healing ligament in rats, 1113
Subject-specific finite element analysis of the human medial collateral ligament during valgus knee loading, 1098

Linearity
Physiological axial compressive preloads increase motion segment stiffness, linearity and hysteresis in all six degrees of freedom for small displacements about the neutral posture, 547

Load sharing
In vivo load sharing among the quadriceps components, 565

Lumbar spine
Effect of compressive follower preload on the flexion–extension response of the human lumbar spine, 540
Physiological axial compressive preloads increase motion segment stiffness, linearity and hysteresis in all six degrees of freedom for small displacements about the neutral posture, 547

Macrophage
Effect of pamidronate on the stimulation of macrophage TNF-α release by ultra-high-molecular-weight polyethylene particles: a role for apoptosis, 81
Hyalgan® has a dose-dependent differential effect on macrophage proliferation and cell death, 744

Maillard reaction
Glucose-mediated in vitro glycation modulates biomechanical integrity of the soft tissues but not hard tissues, 738

MAP kinases
Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts, 1041

Matrix
Type II collagen modulates the composition of extracellular matrix synthesized by articular chondrocytes, 282

MCL
Subject-specific finite element analysis of the human medial collateral ligament during valgus knee loading, 1098

Mechanical characterization
An externally fixed femoral fracture model for mice, 685

Mechanical forces
Gliding properties of the long head of the biceps brachii, 162

Mechanical loading
Bone architecture and disc degeneration in the lumbar spine of mice lacking GDF-8 (myostatin), 1025

Mechanical stimulation
Long-term intermittent shear deformation improves the quality of cartilaginous tissue formed in vitro, 590

Mechanical testing
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224
A reliable externally fixated murine femoral fracture model that accounts for variation in movement between animals, 843

Mechanobiology
Cyclic hydrostatic pressure enhances the chondrogenic phenotype of human mesenchymal progenitor cells differentiated in vitro, 451

Medial collateral ligament
Correlation of healing capacity with vascular response in the anterior cruciate and medial collateral ligaments of the rabbit, 1118
The healing medial collateral ligament following a combined anterior cruciate and medial collateral ligament injury—a biomechanical study in a goat model, 1124
Vascular adaptation of intact joint stabilizing structures in the posterior cruciate ligament deficient rabbit knee, 787

Meniscus
Differential effects of static and dynamic compression on meniscal cell gene expression, 963
The level of compressive load affects conclusions from statistical analyses to determine whether a lateral meniscal autograft restores tibial contact pressure to normal: a study in human cadaveric knees, 459

Mesenchymal progenitor cells
Chondrogenic differentiation of mesenchymal progenitor cells encapsulated in ultrahigh-viscosity alginate, 1090

Mesenchymal stem cell
Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles, 697

Mesenchymal stem cells
Skeletal homeostasis in tissue-engineered bone, 859

Metalloproteinases
IL-1β induces COX2, MMP-1, -3 and -13, ADAMTS-4, IL-1β and IL-6 in human tendon cells, 256

Metastasis
Differences in the cytokine profiles associated with prostate cancer cell induced osteoblastic and osteolytic lesions in bone, 62
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

Microcomputed tomography (μCT)
MicroCT evaluation of normal and osteoarthritic bone structure in human knee specimens, 6

Microparticles
Platelets are mitogenic for periosteum-derived cells, 941

Microscopical and clinical recovery
Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth, 976

Microspheres
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Mineralization
Cryopreservation of tissue engineered constructs for bone, 1005

Miniosmotic pump
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

MMP-1
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

MMP-1, interstitial collagenase
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

MMPs, matrix metalloproteinase
Inhibition of MMP-1 expression by antisense RNA decreases invasiveness of human chondrosarcoma, 1063

Moment distribution
In vivo load sharing among the quadriceps components, 565

Motor control
Simultaneous feedforward recruitment of the vasti in untrained postural tasks can be restored by physical therapy, 553

Mouse
A reliable externally fixated murine femoral fracture model that accounts for variation in movement between animals, 843
GDF-5 deficiency in mice delays Achilles tendon healing, 826

Mouse model
An externally fixed femoral fracture model for mice, 685

Multiple ligament injuries
The healing medial collateral ligament following a combined anterior cruciate and medial collateral ligament injury—a biomechanical study in a goat model, 1124

Muscle atrophy
Early reattachment does not reverse atrophy and fat accumulation of the supraspinatus—an experimental study in rabbits, 386

Muscle forces
THA loading arising from increased femoral anteversion and offset may lead to critical cement stresses, 767

Muscle injuries
Gamma interferon as an antifibrosis agent in skeletal muscle, 798

Muscle mass
Bone architecture and disc degeneration in the lumbar spine of mice lacking GDF-8 (myostatin), 1025

Muscle regeneration
Gamma interferon as an antifibrosis agent in skeletal muscle, 798

Neovascularization
Shock wave therapy induces neovascularization at the tendon–bone junction A study in rabbits, 984

Neuromuscular control
Specificity of muscle action after anterior cruciate ligament injury, 1131

Neuropeptides
Neuronal plasticity in relation to nociception and healing of rat achilles tendon, 432

Nitric oxide
Nitric oxide inhibition of IGF-1 stimulated proteoglycan synthesis: role of cGMP, 914

NMR
Determination of cortical bone porosity and pore size distribution using a low field pulsed NMR approach, 312

Non-enzymatic glycation
Glucose-mediated in vitro glycation modulates biomechanical integrity of the soft tissues but not hard tissues, 738

Non-traumatic rabbit osteonecrosis model
Evaluation of femoral perfusion in a non-traumatic rabbit osteonecrosis model with T2*-weighted dynamic MRI, 341

Noninvasive fracture model
Noninvasive fatigue fracture model of the rat ulna, 1018

Nonunion
Development of an atrophic nonunion model and comparison to a closed healing fracture in rat femur, 503

NSAIDs
Differential inhibition of fracture healing by non-selective and cyclooxygenase-2 selective non-steroidal anti-inflammatory drugs, 670

OP-1, osteo-protein-1 or bone morphogenic protein-7
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

OPG
RANKL is an essential cytokine mediator of polymethylmethacrylate particle-induced osteoclastogenesis, 202

Opioids
Purification and quantification of opioid peptides in bone and joint tissues—a methodological study in the rat, 465

Optical tweezers
Temporal effects of cell adhesion on mechanical characteristics of the single chondrocyte, 88

Osseointegration
Autologous osteoblasts enhance osseointegration of porous titanium implants, 213

Osteoarthritis
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224
Alterations in endogenous osteogenic protein-1 with degeneration of human articular cartilage, 899
Bone architecture and disc degeneration in the lumbar spine of mice lacking GDF-8 (myostatin), 1025
Fibronectin metabolism of cartilage explants in response to the frequency of intermittent loading, 1081
Hyalgan® has a dose-dependent differential effect on macrophage proliferation and cell death, 744
MicroCT evaluation of normal and osteoarthritic bone structure in human knee specimens, 6
Quadriceps strength and volitional activation before and after total knee arthroplasty for osteoarthritis, 775
Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes, 96
Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis, 792
Spontaneous and experimental osteoarthritis in dog: similarities and differences in proteoglycan levels, 730

Osteoblast
Effects of receptor activator of NFκB (RANK) signaling blockade on fracture healing, 676

Osteoblastic lesions
Differences in the cytokine profiles associated with prostate cancer cell induced osteoblastic and osteolytic lesions in bone, 62

Osteoblasts
Pretreatment of bone with osteoclasts affects phenotypic expression of osteoblast-like cells, 638
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Osteoclast
Bone loss following tendon laceration, repair and passive mobilization, 990
Effects of receptor activator of NFκB (RANK) signaling blockade on fracture healing, 676

Osteoclast precursor cells
RANKL is an essential cytokine mediator of polymethylmethacrylate particle-induced osteoclastogenesis, 202

Osteoclastogenesis
RANKL is an essential cytokine mediator of polymethylmethacrylate particle-induced osteoclastogenesis, 202

Osteoclasts
Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts, 1041
Pretreatment of bone with osteoclasts affects phenotypic expression of osteoblast-like cells, 638

Osteocytes
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Osteogenic protein-1
Alterations in endogenous osteogenic protein-1 with degeneration of human articular cartilage, 899
Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads, 922

Osteoinductivity
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Osteon
Osteon pullout in the equine third metacarpal bone: effects of ex vivo fatigue, 481

Osteonecrosis
Measurement of lesion area and volume by three-dimensional spoiled gradient-echo MR imaging in osteonecrosis of the femoral head, 850

Osteopetrosis
Effects of receptor activator of NFκB (RANK) signaling blockade on fracture healing, 676

Osteoporosis
Biomechanical evaluation of healing in a non-critical defect in a large animal model of osteoporosis, 836

Osteoprotegerin
Differences in the cytokine profiles associated with prostate cancer cell induced osteoblastic and osteolytic lesions in bone, 62

Osteosarcoma
Chemotherapeutic agents sensitize sarcoma cell lines to tumor necrosis factor-related apoptosis-inducing ligand-induced caspase-8 activation, apoptosis and loss of mitochondrial membrane potential, 949

Osteotomy
rhBMP-2 injected in a calcium phosphate paste (α-BSM) accelerates healing in the rabbit ulnar osteotomy model, 997

Oxygen
Effects of transcutaneous topical injection of oxygen on vascular endothelial growth factor gene into the healing ligament in rats, 1113

p38
Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts, 1041

Pain
Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis, 792

Pain measurement
Neuronal plasticity in relation to nociception and healing of rat achilles tendon, 432

Parathyroid hormone
Skeletal homeostasis in tissue-engineered bone, 859

Parathyroid hormone-related peptide
Expression of parathyroid hormone-related peptide and insulin-like growth factor I during rat fracture healing, 511

Patella
Quantitative measurement of patellofemoral joint stability: force–displacement behavior of the human patella in vitro, 780

Patellofemoral joint
Quantitative measurement of patellofemoral joint stability: force–displacement behavior of the human patella in vitro, 780

PCNA
Shock wave therapy induces neovascularization at the tendon–bone junction A study in rabbits, 984

PCR
Detection of chimerism following vascularized bone allotransplantation by polymerase chain reaction using a Y-chromosome specific primer, 1056

Perichondrium
Adhesion of perichondrial cells to a polylactic acid scaffold, 584

Periosteum
Induction of CD-RAP mRNA during periosteal chondrogenesis, 296

Peripheral nervous system
Neuronal plasticity in relation to nociception and healing of rat achilles tendon, 432

Physical stimulation
Physiologic weight-bearing increases new vessel formation during distraction osteogenesis: a micro-tomographic imaging study, 489

Pivot shift test
A quantitative analysis of valgus torque on the ACL: a human cadaveric study, 1107

Plantar fasciitis
Extracorporeal shock wave therapy for plantar fasciitis. A double blind randomised controlled trial, 937

Platelet-derived growth factor
Platelets are mitogenic for periosteum-derived cells, 941

Platelet-rich plasma
Platelets are mitogenic for periosteum-derived cells, 941

PMMA
Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts, 1041

Polyethylene particles
Effect of pamidronate on the stimulation of macrophage TNF-α release by ultra-high-molecular-weight polyethylene particles: a role for apoptosis, 81

Popliteus
The anatomy of the posterolateral aspect of the rabbit knee, 723

Pore size
Determination of cortical bone porosity and pore size distribution using a low field pulsed NMR approach, 312

Porous coated implant
The effects of sex and estrogen therapy on bone ingrowth into porous coated implant, 1033

Posterolateral corner
The anatomy of the posterolateral aspect of the rabbit knee, 723

Postural perturbation
Simultaneous feedforward recruitment of the vasti in untrained postural tasks can be restored by physical therapy, 553

Proliferation
Hyalgan® has a dose-dependent differential effect on macrophage proliferation and cell death, 744

Proprioception
Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis, 792

Prospective
Quadriceps strength and volitional activation before and after total knee arthroplasty for osteoarthritis, 775

Prostaglandin E_2
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Prostate cancer
Differences in the cytokine profiles associated with prostate cancer cell induced osteoblastic and osteolytic lesions in bone, 62

Proteoglycan
Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads, 922

Pulsed electromagnetic fields
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

Quadriceps
In vivo load sharing among the quadriceps components, 565
Quadriceps strength and volitional activation before and after total knee arthroplasty for osteoarthritis, 775
Simultaneous feedforward recruitment of the vasti in untrained postural tasks can be restored by physical therapy, 553

Quadriceps femoris muscle setting
Electromyographic analysis of a modified maneuver for quadriceps femoris muscle setting with co-contraction of the hamstrings, 559

Rabbit
Effect of dynamic hydrostatic pressure on rabbit intervertebral disc cells, 597
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335
Evaluation of collagen fiber maturation and ordering in regenerating tendons employing H-1 double quantum filtered NMR spectroscopy, 149

Radioactive microspheres
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Radiofrequency energy
A comparison of joint stability between anterior cruciate intact and deficient knees: a new canine model of anterior cruciate ligament disruption, 224

Radioimmunoassay
Purification and quantification of opioid peptides in bone and joint tissues—a methodological study in the rat, 465

RANK
Effects of receptor activator of NFκB (RANK) signaling blockade on fracture healing, 676

RANKL
RANKL is an essential cytokine mediator of polymethylmethacrylate particle-induced osteoclastogenesis, 202

Rat
Influence of a standardized closed soft tissue trauma on resistance to local infection. An experimental study in rats, 373

Rats
Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth, 976
Physiologic weight-bearing increases new vessel formation during distraction osteogenesis: a micro-tomographic imaging study, 489

Real-time PCR
Bilaterally increased VEGF-levels in muscles during experimental unilateral callus distraction, 805

Regeneration
Neuronal plasticity in relation to nociception and healing of rat achilles tendon, 432

Remodeling
Noninvasive fatigue fracture model of the rat ulna, 1018
Osteon pullout in the equine third metacarpal bone: effects of ex vivo fatigue, 481

Repetitive loads
Time, stress, and location dependent chondrocyte death and collagen damage in cyclically loaded articular cartilage, 888

Reply to Letter to the Editor
961

Robotics/universal force–moment sensor (UFS) testing system
A quantitative analysis of valgus torque on the ACL: a human cadaveric study, 1107

Rotator cuff suture
Early reattachment does not reverse atrophy and fat accumulation of the supraspinatus—an experimental study in rabbits, 386

Rotator cuff tear
Early reattachment does not reverse atrophy and fat accumulation of the supraspinatus—an experimental study in rabbits, 386

RT-PCR
Induction of CD-RAP mRNA during periosteal chondrogenesis, 296

SDS, sodium dodecyl sulfate
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819

Sensitization
Chemotherapeutic agents sensitize sarcoma cell lines to tumor necrosis factor-related apoptosis-inducing ligand-induced caspase-8 activation, apoptosis and loss of mitochondrial membrane potential, 949

SGF, stem cell growth factor
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

Shear
Long-term intermittent shear deformation improves the quality of cartilaginous tissue formed in vitro, 590

Sheep
Biomechanical evaluation of healing in a non-critical defect in a large animal model of osteoporosis, 836

Shock wave
Extracorporeal shock wave therapy for plantar fasciitis. A double blind randomised controlled trial, 937

Shock wave therapy
Shock wave therapy induces neovascularization at the tendon–bone junction A study in rabbits, 984

Shoulder
Variation of biomechanical, structural, and compositional properties along the tendon to bone insertion site, 413

Signal transduction
Epigallocatechin-3-gallate selectively inhibits interleukin-1β-induced activation of mitogen activated protein kinase subgroup c-*Jun N*-terminal kinase in human osteoarthritis chondrocytes, 102

Signaling
Type II collagen modulates the composition of extracellular matrix synthesized by articular chondrocytes, 282

Smad6
Smad6 is induced by BMP-2 and modulates chondrocyte differentiation, 908

Small proteoglycans
Spontaneous and experimental osteoarthritis in dog: similarities and differences in proteoglycan levels, 730

Sprague-Dawley rats
Effects of nicotine on bone mass and strength in aged female rats, 14

Stem cell
Cyclic hydrostatic pressure enhances the chondrogenic phenotype of human mesenchymal progenitor cells differentiated in vitro, 451

Stiffness
Physiological axial compressive preloads increase motion segment stiffness, linearity and hysteresis in all six degrees of freedom for small displacements about the neutral posture, 547

Stimulation
Shear movement at the fracture site delays healing in a diaphyseal fracture model, 1011

Strength
The relationship between knee strength and functional stability before and after anterior cruciate ligament reconstruction, 231

Structural integrity
Glucose-mediated in vitro glycation modulates biomechanical integrity of the soft tissues but not hard tissues, 738

Subject-specific modeling
Subject-specific finite element analysis of the human medial collateral ligament during valgus knee loading, 1098

Subpopulations
The use of specific chondrocyte populations to modulate the properties of tissue-engineered cartilage, 132

Superficial tangential zone
Time, stress, and location dependent chondrocyte death and collagen damage in cyclically loaded articular cartilage, 888

Surface
Cell death after cartilage impact occurs around matrix cracks, 881

Surface morphology
Pretreatment of bone with osteoclasts affects phenotypic expression of osteoblast-like cells, 638

T_2 relaxation
Determination of cortical bone porosity and pore size distribution using a low field pulsed NMR approach, 312

T2*-weighted dynamic MRI
Evaluation of femoral perfusion in a non-traumatic rabbit osteonecrosis model with T2*-weighted dynamic MRI, 341

Tendon
IL-1β induces COX2, MMP-1, -3 and -13, ADAMTS-4, IL-1β and IL-6 in human tendon cells, 256
Inhibition of cell migration from tendon explants into fibrin clots by extracts derived from cheese whey is largely due to transforming growth factor-beta, 157
Tendon cells produce gelatinases in response to type I collagen attachment, 442
Variation of biomechanical, structural, and compositional properties along the tendon to bone insertion site, 413

Tendon healing
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819
Inhibition of tendon cell migration by dexamethasone is correlated with reduced alpha-smooth muscle actin gene expression: a potential mechanism of delayed tendon healing, 265

Tendon laceration
Bone loss following tendon laceration, repair and passive mobilization, 990

Tendon repair
Bone loss following tendon laceration, repair and passive mobilization, 990
GDF-5 deficiency in mice delays Achilles tendon healing, 826

Tendon rupture
Neuronal plasticity in relation to nociception and healing of rat achilles tendon, 432

Tendon–bone junction
Shock wave therapy induces neovascularization at the tendon–bone junction A study in rabbits, 984

TGF-β, transforming growth factor-beta
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

TGF-β1
Type II collagen modulates the composition of extracellular matrix synthesized by articular chondrocytes, 282

Therapeutic interventions
The use of intra-articular Na-hyaluronate as a potential chondroprotective device in experimentally induced acute articular cartilage injury and repair in rabbits, 305

Three-dimensional spoiled gradient-echo (3D SPGR) imaging
Measurement of lesion area and volume by three-dimensional spoiled gradient-echo MR imaging in osteonecrosis of the femoral head, 850

Tibia
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Tissue biomechanics
Glucose-mediated in vitro glycation modulates biomechanical integrity of the soft tissues but not hard tissues, 738

Tissue engineering
Adhesion of perichondrial cells to a polylactic acid scaffold, 584
Autologous osteoblasts enhance osseointegration of porous titanium implants, 213
Bone induction by *BMP-2* transduced stem cells derived from human fat, 622
Cryopreservation of tissue engineered constructs for bone, 1005
Skeletal homeostasis in tissue-engineered bone, 859
The use of specific chondrocyte populations to modulate the properties of tissue-engineered cartilage, 132

Tissue-engineering
Long-term intermittent shear deformation improves the quality of cartilaginous tissue formed in vitro, 590

Titanium
Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles, 697

Titanium implants
Autologous osteoblasts enhance osseointegration of porous titanium implants, 213

Tobacco-smoking
Effects of nicotine on bone mass and strength in aged female rats, 14

Torsional strength
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Total hip arthroplasty
Bone mineral density and biochemical markers of bone turnover in aseptic loosening after total hip arthroplasty, 691
THA loading arising from increased femoral anteversion and offset may lead to critical cement stresses, 767

Total hip replacements
Four-year study of cobalt and chromium blood levels in patients managed with two different metal-on-metal total hip replacements, 189

Toxic effect
The effects of therapeutic vs. high-intensity ultrasound on the rabbit growth plate, 865

Trabecular bone
Bone architecture and disc degeneration in the lumbar spine of mice lacking GDF-8 (myostatin), 1025

Trabecular structure
MicroCT evaluation of normal and osteoarthritic bone structure in human knee specimens, 6

TRAIL
Chemotherapeutic agents sensitize sarcoma cell lines to tumor necrosis factor-related apoptosis-inducing ligand-induced caspase-8 activation, apoptosis and loss of mitochondrial membrane potential, 949

Training
Simultaneous feedforward recruitment of the vasti in untrained postural tasks can be restored by physical therapy, 553

Transcutaneous topical injection
Effects of transcutaneous topical injection of oxygen on vascular endothelial growth factor gene into the healing ligament in rats, 1113

Transforming growth factor β1
Gamma interferon as an antifibrosis agent in skeletal muscle, 798

Transforming growth factor-β
Inhibition of cell migration from tendon explants into fibrin clots by extracts derived from cheese whey is largely due to transforming growth factor-beta, 157

Transforming growth factor-β1
Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast-like cells, 326

TRAP
Rapid new bone tissue remodeling during distraction osteogenesis is associated with apoptosis, 28

Treatment
Extracorporeal shock wave therapy for plantar fasciitis. A double blind randomised controlled trial, 937

TUNEL
Rapid new bone tissue remodeling during distraction osteogenesis is associated with apoptosis, 28
The fate of soft callus chondrocytes during long bone fracture repair, 54

Type I collagen
Tendon cells produce gelatinases in response to type I collagen attachment, 442

Type X collagen
Smad6 is induced by BMP-2 and modulates chondrocyte differentiation, 908

Ultrasound
The effects of therapeutic vs. high-intensity ultrasound on the rabbit growth plate, 865

uPA, urokinase-type plasminogen activator
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819

uPAR, urokinase-type plasminogen activator receptor
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819

Urokinase
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819

Urokinase receptor
Expression of urokinase-type plasminogen activator and its receptor is up-regulated during tendon healing, 819

Valgus torque
A quantitative analysis of valgus torque on the ACL: a human cadaveric study, 1107

Vascular endothelial growth factor
Effects of transcutaneous topical injection of oxygen on vascular endothelial growth factor gene into the healing ligament in rats, 1113

Vascular endothelial growth factor (VEGF)
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

VEGF
Bilaterally increased VEGF-levels in muscles during experimental unilateral callus distraction, 805
Shock wave therapy induces neovascularization at the tendon–bone junction
A study in rabbits, 984

VEGF, vascular endothelial growth factor
Quantitative and sensitive in vitro assay for osteoinductive activity of demineralized bone matrix, 648

VEGF-inhibitor
Effects of locally applied vascular endothelial growth factor (VEGF) and VEGF-inhibitor to the rabbit tibia during distraction osteogenesis, 335

Viscoelastic properties
Variation of biomechanical, structural, and compositional properties along the tendon to bone insertion site, 413

Viscosupplementation
The use of intra-articular Na-hyaluronate as a potential chondroprotective device in experimentally induced acute articular cartilage injury and repair in rabbits, 305

Volume
Measurement of lesion area and volume by three-dimensional spoiled gradient-echo MR imaging in osteonecrosis of the femoral head, 850

Wear particles
Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles, 697

Whey
Inhibition of cell migration from tendon explants into fibrin clots by extracts derived from cheese whey is largely due to transforming growth factor-beta, 157

Wound healing
Effects of transcutaneous topical injection of oxygen on vascular endothelial growth factor gene into the healing ligament in rats, 1113

X-ray
The effects of therapeutic vs. high-intensity ultrasound on the rabbit growth plate, 865

Y-chromosome
Detection of chimerism following vascularized bone allotransplantation by polymerase chain reaction using a Y-chromosome specific primer, 1056

Zirconia
Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles, 697